

## Nachzuchtstatistik der Unterfamilie 'Loriinae - Eigentliche Loris' (2007)

| Jahr | Deutscher Name                           | Wissensch. Name                                                   | Paare | 1,0 | 0,1 | Unbek. | Gesamt |
|------|------------------------------------------|-------------------------------------------------------------------|-------|-----|-----|--------|--------|
| 2007 | BRAUNLORI                                | Chalcopsitta duivenbodei duivenbodei (Dubois, 1884)               | 3     | 2   | 2   | 0      | 4      |
| 2007 | JOSEPHINENLORI                           | Charmosyna josefinae josefinae (Finsch, 1873)                     | 6     | 5   | 10  | 0      | 15     |
| 2007 | SEPIK-JOSEPHINENLORI                     | Charmosyna josefinae sepikiana (Neumann, 1922)                    | 0     | 0   | 0   | 3      | 3      |
| 2007 | STELLAS PAPUALORI                        | Charmosyna papou stellae (A.B. Meyer, 1886)                       | 8     | 5   | 14  | 0      | 19     |
| 2007 | HARTERTS-GOLDSTRICHELLORI                | Charmosyna pulchella rothschildi (Hartert, 1930)                  | 1     | 0   | 5   | 0      | 5      |
| 2007 | AMBOINA-ROTLORI                          | Eos bornea bornea (Linné, 1758)                                   | 1     | 0   | 0   | 5      | 5      |
| 2007 | STRICHELLORI (BLAUSTRICHELLORI)          | Eos reticulata (S. Müller, 1841)                                  | 1     | 0   | 0   | 2      | 2      |
| 2007 | OBI KAPUZENLORI                          | Eos squamata obiensis (Rothschild, 1899)                          | 2     | 1   | 1   | 0      | 2      |
| 2007 | MOSCHUSLORI                              | Glossopsitta concinna (Shaw, 1791)                                | 3     | 4   | 2   | 1      | 7      |
| 2007 | BLAUBÜRZELIGER SCHÖNLORI                 | Hypocharmosyna placentis ornata (Mayr, 1940)                      | 3     | 3   | 3   | 0      | 6      |
| 2007 | ROTBÜRZELLORI (ROTSTIRNLORI)             | Hypocharmosyna rubronotata rubronotata (Wallace, 1862)            | 1     | 1   | 0   | 0      | 1      |
| 2007 | GRÜNSCHWANZLORI                          | Lorius chlorocercus (Gould, 1856)                                 | 6     | 9   | 9   | 0      | 18     |
| 2007 | ERZLORI                                  | Lorius domicellus (Linné, 1758)                                   | 3     | 2   | 2   | 0      | 4      |
| 2007 | GELBMANTELLORI                           | Lorius garrulus garullus (Linné, 1758)                            | 2     | 0   | 1   | 6      | 7      |
| 2007 | FRAUENLORI                               | Lorius lory lory (Linné, 1758)                                    | 7     | 3   | 3   | 1      | 7      |
| 2007 | ARFAK-BERGZIERLORI (ARFAKLORI)           | Oreopsittacus arfaki arfaki (A.B. Meyer, 1874)                    | 2     | 1   | 2   | 0      | 3      |
| 2007 | GROSSER BERGZIERLORI                     | Oreopsittacus arfaki major (Ogilvie-Grant, 1914)                  | 8     | 8   | 7   | 0      | 15     |
| 2007 | EINSIEDLERLORI (KRAGENLORI)              | Phigys solitarius (Suckow, 1800)                                  | 1     | 3   | 1   | 0      | 4      |
| 2007 | VEILCHENLORI                             | Psitteuteles goldiei (Sharpe, 1882)                               | 3     | 3   | 3   | 0      | 6      |
| 2007 | BUNTLORI                                 | Psitteuteles versicolor (Lear, 1831)                              | 1     | 1   | 1   | 0      | 2      |
| 2007 | GELBKOPFLORI                             | Trichoglossus euteles (Temminck, 1835)                            | 2     | 1   | 3   | 0      | 4      |
| 2007 | FORSTENLORI                              | Trichoglossus haematodus forsteni (Bonaparte, 1850)               | 2     | 0   | 1   | 2      | 3      |
| 2007 | BREITBINDEN-ALLFARBLORI                  | Trichoglossus haematodus haematodus (Linné, 1771)                 | 6     | 3   | 2   | 5      | 10     |
| 2007 | MASSENA-ALLFARBLORI (SCHMALBINDENLORI)   | Trichoglossus haematodus massena (Bonaparte, 1854)                | 1     | 1   | 1   | 0      | 2      |
| 2007 | MITCHELL-ALLFARBLORI                     | Trichoglossus haematodus mitchellii (G.R. Gray, 1859)             | 3     | 3   | 4   | 0      | 7      |
| 2007 | GEBIRGSLORI (LORI VON DEN BLAUEN BERGEN) | Trichoglossus haematodus moluccanus (Gmelin 1788)                 | 36    | 37  | 27  | 46     | 110    |
| 2007 | ROSENBERGS-ALLFARBLORI                   | Trichoglossus haematodus rosenbergii (Schlegel, 1871)             | 3     | 6   | 4   | 0      | 10     |
| 2007 | ROTNACKENLORI                            | Trichoglossus haematodus rubritorquis (Vigors u. Horsfield, 1826) | 6     | 12  | 13  | 1      | 26     |
| 2007 | WEBER-ALLFARBLORI                        | Trichoglossus haematodus weberi (Hartert, 1898)                   | 2     | 1   | 1   | 1      | 3      |
| 2007 | SCHMUCKLORI                              | Trichoglossus ornatus (Linné, 1758)                               | 5     | 5   | 10  | 0      | 15     |
| 2007 | BLAUKÄPPCHEN (BLAUKAPPENLORI)            | Vini australis (Gmelin, 1788)                                     | 4     | 4   | 3   | 0      | 7      |